

Title: Impact of a 1755-like tsunami in Huelva, Spain

Author(s): Lima, V. V. ^[1]; Miranda, J. M. ^[1]; **Baptista, M. A.** ^[1,2]; Catalao, J. ^[1]; Gonzalez, M. ^[3]; Otero, L. ^[3]; Olabarrieta, M. ^[3]; Alvarez-Gomez, J. A. ^[4]; Carreno, E. ^[4]

Source: Natural Hazards and Earth Systems Sciences

Volume: 10 **Issue:** 1 **Pages:** 139-148 **Published:** 2010

Document Type: Article

Language: English

Abstract: Coastal areas are highly exposed to natural hazards associated with the sea. In all cases where there is historical evidence for devastating tsunamis, as is the case of the southern coasts of the Iberian Peninsula, there is a need for quantitative hazard tsunami assessment to support spatial planning. Also, local authorities must be able to act towards the population protection in a preemptive way, to inform 'what to do' and 'where to go' and in an alarm, to make people aware of the incoming danger. With this in mind, we investigated the inundation extent, run-up and water depths, of a 1755-like event on the region of Huelva, located on the Spanish southwestern coast, one of the regions that was affected in the past by several high energy events, as proved by historical documents and sedimentological data. Modelling was made with a slightly modified version of the COMCOT (Cornell Multi-grid Coupled Tsunami Model) code. Sensitivity tests were performed for a single source in order to understand the relevance and influence of the source parameters in the inundation extent and the fundamental impact parameters. We show that a 1755-like event will have a dramatic impact in a large area close to Huelva inundating an area between 82 and 92 km(2) and reaching maximum run-up around 5 m. In this sense our results show that small variations on the characteristics of the tsunami source are not too significant for the impact assessment. We show that the maximum flow depth and the maximum run-up increase with the average slip on the source, while the strike of the fault is not a critical factor as Huelva is significantly far away from the potential sources identified up to now. We also show that the maximum flow depth within the inundated area is very dependent on the tidal level, while maximum run-up is less affected, as a consequence of the complex morphology of the area.

KeyWords Plus: 1755 Lisbon Tsunami; Numerical Simulations; Earthquake; Record

Reprint Address: Lima, VV (reprint author) - Univ Lisbon, IDL, Campo Grande, Ed C8, P-1749016 Lisbon, Portugal

Addresses:

[1] Univ Lisbon, IDL, P-1749016 Lisbon, Portugal

[2] IPL, Inst Super Engn Lisboa, Lisbon, Portugal

[3] Univ Cantabria, IH Cantabria, Environm Hydraul Inst, ETSI Caminos Canales & Puertos, E-39005 Santander, Spain

[4] Ctr Nacl Informac Geog, Madrid 28003, Spain

E-mail Addresses: vcvalima@gmail.com

Funding:

Funding Agency	Grant Number
TRANSFER	STREP 37058
NEAREST	STREP 037110

Publisher: Copernicus Gesellschaft MBH

Publisher Address: Bahnhofsallee 1E, Gottingen, 37081, Germany

ISSN: 1561-8633

Citation: LIMA, V. V.; MIRANDA, J. M.; BAPTISTA, M. A.; CATALAO, J.; GONZALEZ, M.; OTERO, L.; OLABARRIETA, M.; ALVAREZ-GOMEZ, J. A.; CARRENO, E. - Impact of a 1755-like tsunami in Huelva, Spain. Natural Hazards and Earth Systems Sciences. ISSN 1561-8633. Vol. 10, nr. 1 (2010), p. 139-148.